

ON THE EXISTENCE OF ROTATED D_n -LATTICES CONSTRUCTED VIA GALOIS EXTENSIONS

GRASIELE C. JORGE AND SUELI I. R. COSTA

IMECC-UNICAMP ,

651, St. Sérgio Buarque de Holanda, 13083-859, Campinas, SP, BRAZIL

Emails: [grajorge, sueli] @ime.unicamp.br

ABSTRACT. In this paper we construct families of rotated D_n -lattices which may be suitable for signal transmission over both Gaussian and Rayleigh fading channels via subfields of cyclotomic fields. These constructions exhibit full diversity and good minimum product distance which are important parameters related to the signal transmission error probability. It is also shown that for some Galois extensions $\mathbb{K}|\mathbb{Q}$ it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}}$.

Mathematical subject classification: 11H06 - 11R80 - 97N70

Keywords: D_n -lattices, Galois extensions, Full diversity lattices

1. INTRODUCTION

A *lattice* Λ is a discrete additive subgroup of $\mathbb{R}^n$. Equivalently, $\Lambda \subseteq \mathbb{R}^n$ is a lattice iff there are linearly independent vectors $\mathbf{v}_1, \dots, \mathbf{v}_m \in \mathbb{R}^n$, $m \leq n$, such that any $\mathbf{y} \in \Lambda$ can be written as $\mathbf{y} = \sum_{i=1}^m \alpha_i \mathbf{v}_i$, $\alpha_i \in \mathbb{Z}$. The set $\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$ is called a *basis* for Λ . A matrix M whose rows are these vectors is said to be a *generator matrix* for Λ while the associated *Gram matrix* is $G = MM^t$. The *determinant* of Λ is $\det \Lambda = \det G$ and it is an invariant under basis change [3].

Lattices are used in applications of different areas, particularly in information theory and more recently in cryptography. In this paper we attempt to construct lattices with full rank ($m=n$) which may be suitable for signal transmission over both Gaussian and Rayleigh fading channels [2]. For this purpose the lattice parameters that we consider here are packing density, diversity and minimum product distance [3, 5].

The *packing density*, $\Delta(\Lambda)$, of a lattice Λ is the proportion of the space $\mathbb{R}^n$ covered by the union of congruent disjoint spheres of maximum radius centered at the points of Λ [3]. A lattice Λ has *diversity* $k \leq n$ if k is the maximum number such that for all $\mathbf{y} = (y_1, \dots, y_n) \in \Lambda$, $\mathbf{y} \neq \mathbf{0}$, there are at least k nonzero coordinates. Given a

This work was partially supported by CAPES 2548/2010, CNPq 150802/2012-9, 309561/2009-4 and FAPESP 2007/56052-8.

full rank lattice with full diversity $\Lambda \subseteq \mathbb{R}^n$ ($k = n$) the *minimum product distance* is defined as $d_{\min}(\Lambda) = \min \{ \prod_{i=1}^n |y_i|, \mathbf{y} = (y_1, \dots, y_n) \in \Lambda, \mathbf{y} \neq \mathbf{0} \}$ [5].

Special algebraic lattice constructions allows to derive certain lattice parameters, which are usually difficult to be calculated for general lattices in $\mathbb{R}^n$, such as the packing density and the minimum product distance. Some of these constructions can be found in [2, 5–8].

In [9] we have constructed families of full diversity rotated D_n -lattices through of a twisted homomorphism and a free $\mathbb{Z}$ -module of rank n contained in the totally real subfield $\mathbb{Q}(\zeta_m + \zeta_m^{-1})$ of the cyclotomic field $\mathbb{Q}(\zeta_m)$ for $n = \phi(m)/2$, where ϕ is the Euler function. For $\mathbb{K}_1 = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $r \geq 5$, rotated D_{2^r-2} -lattices were obtained via free $\mathbb{Z}$ -modules which are ideals of $\mathcal{O}_{\mathbb{K}_1}$ whereas for $\mathbb{K}_2 = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$, $p \geq 7$ prime, rotated $D_{(p-1)/2}$ -lattices were constructed via free $\mathbb{Z}$ -modules $M \subseteq \mathcal{O}_{\mathbb{K}_2}$ of rank $n = [\mathbb{K}_2 : \mathbb{Q}]$ that are not ideals of $\mathcal{O}_{\mathbb{K}_2}$.

In this paper, considering the compositum $\mathbb{K}_3$ of $\mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$ and $\mathbb{Q}(\zeta_p + \zeta_p^{-1})$ and the compositum $\mathbb{K}_4$ of $\mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})$ and $\mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})$, where p, p_1, p_2 are prime numbers, $p_1 \neq p_2$, we construct families of rotated D_n -lattices via free $\mathbb{Z}$ -modules of rank n that are not ideals. In both cases we get a closed-form for their minimum product distances. If it was possible to construct such rotated D_n -lattices via principal ideals of $\mathcal{O}_{\mathbb{K}_i}$, $i = 2, 3, 4$, their minimum product distances should be twice the ones obtained in our construction and therefore more appropriate to our purpose. However, we show here that in the cases we have considered it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}_i}$, $i = 2, 3, 4$. Moreover, we present a necessary condition to construct these lattices via a Galois extension $\mathbb{K}|\mathbb{Q}$. These results are related to an open problem stated in [4]: Given a lattice Λ , is there a number field $\mathbb{K}$ such that it is possible to construct Λ via $\mathcal{O}_{\mathbb{K}}$?

This paper is organized as follows. In Section 2 we collect some definitions and results on ideal lattices. In Section 3 new constructions of rotated D_n -lattices obtained via the compositum of some subfields of cyclotomic fields are derived. Section 4 is devoted to a discussion on the existence of rotated D_n -lattices constructed via fractional ideals of a Galois extension $\mathbb{K}|\mathbb{Q}$.

2. IDEAL LATTICES

The construction of ideal lattices presented here was introduced in [4].

In what follows we use some classical notation for number fields [11–13]. Let $\mathbb{K}$ be a Galois extension of degree $n = [\mathbb{K} : \mathbb{Q}]$, $\text{Gal}(\mathbb{K}|\mathbb{Q}) = \{\sigma_i\}_{i=1}^n$ the Galois group, $d_{\mathbb{K}}$ the discriminant of $\mathbb{K}|\mathbb{Q}$, $\mathcal{O}_{\mathbb{K}}$ its ring of integers, $N_{\mathbb{K}|\mathbb{Q}}(x)$ and $\text{Tr}_{\mathbb{K}|\mathbb{Q}}(x)$ the norm and the trace of an element $x \in \mathbb{K}$, respectively.

Definition 2.1. Let $\mathbb{K}$ be a totally real number field and $\alpha \in \mathbb{K}$ such that $\alpha_i = \sigma_i(\alpha) > 0$ for all $i = 1, \dots, n$. The homomorphism

$$\begin{aligned} \sigma_\alpha : \mathbb{K} &\longrightarrow \mathbb{R}^n \\ x &\longmapsto (\sqrt{\alpha_1}\sigma_1(x), \dots, \sqrt{\alpha_n}\sigma_n(x)) \end{aligned}$$

is called *twisted homomorphism*. When $\alpha = 1$ the twisted homomorphism is the so called *Minkowski homomorphism*.

Proposition 2.2. [5] If $\mathcal{I} \subseteq \mathbb{K}$ is a free $\mathbb{Z}$ -module of rank n with $\mathbb{Z}$ -basis $\{w_1, \dots, w_n\}$, then the image $\Lambda = \sigma_\alpha(\mathcal{I})$ is a lattice in $\mathbb{R}^n$ with basis $\{\sigma_\alpha(w_1), \dots, \sigma_\alpha(w_n)\}$. Moreover, if $\mathbb{K}$ is totally real, Λ is a full diversity lattice which has $G = M.M^t = (Tr_{\mathbb{K}|\mathbb{Q}}(\alpha w_i w_j))_{i,j=1}^n$ as a Gram matrix.

Proposition 2.3. [9] Let $\mathbb{K}$ be a totally real field number with $[\mathbb{K} : \mathbb{Q}] = n$ and $\mathcal{I} \subseteq \mathbb{K}$ a free $\mathbb{Z}$ -module of rank n . The minimum product distance of $\Lambda = \sigma_\alpha(\mathcal{I})$ is $d_{p,min}(\Lambda) = \sqrt{N_{\mathbb{K}|\mathbb{Q}}(\alpha)} \min_{0 \neq y \in \mathcal{I}} |N_{\mathbb{K}|\mathbb{Q}}(y)|$.

Proposition 2.4. [5] If $\mathbb{K}$ is a totally real field number, $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is a principal ideal and $\Lambda = \sigma_\alpha(\mathcal{I})$, then $d_{p,min}(\Lambda) = \sqrt{\frac{\det(\Lambda)}{|d_{\mathbb{K}}|}}$.

Definition 2.5. The *relative minimum product distance* of Λ , denoted by $\mathbf{d}_{p,rel}(\Lambda)$, is the minimum product distance of a scaled version of Λ with unitary minimum norm vector.

Proposition 2.7 was presented in [4] in the case in which $\mathcal{I} \subseteq \mathbb{K}$ is a fractional ideal. We can extend it to free $\mathbb{Z}$ -modules of rank n . Using the next lemma the proof of Proposition 2.7 is straightforward.

Lemma 2.6. If $\mathcal{I} \subseteq \mathbb{K}$ is a free $\mathbb{Z}$ -module of rank n , then there is $d \in \mathbb{Z} - \{0\}$ such that $d\mathcal{I} \subset \mathcal{O}_{\mathbb{K}}$.

Proof. Let $\alpha \in \mathbb{K}$ such that $\mathbb{K} = \mathbb{Q}(\alpha)$ and $\{1, \alpha, \dots, \alpha^{n-1}\}$ is a basis of $\mathbb{K}$ over $\mathbb{Q}$. Let $\{\gamma_1, \dots, \gamma_n\}$ be a $\mathbb{Z}$ -basis of $\mathcal{I}$. For any i , $\gamma_i = \sum_{j=0}^{n-1} a_{ij} \alpha^j$ where $a_{ij} \in \mathbb{Q}$, for all $i = 1, \dots, n$ and $j = 0, 1, \dots, n-1$. Since $a_{ij} \in \mathbb{Q}$ for all i, j , it follows that $a_{ij} = \frac{b_{ij}}{c_{ij}}$ with $b_{ij}, c_{ij} \in \mathbb{Z}$ and $c_{ij} \neq 0$ for all i, j . If $d = \text{lcm}\{c_{ij}; i = 1, \dots, n, j = 0, 1, \dots, n-1\}$, then $d\gamma_i \in \mathbb{Z}[\alpha]$ for all $i = 1, \dots, n$, what implies $d\mathcal{I} = d \sum_{i=1}^n \mathbb{Z}\gamma_i = \sum_{i=1}^n \mathbb{Z}d\gamma_i \subset \mathbb{Z}[\alpha] \subset \mathcal{O}_{\mathbb{K}}$. $\square$

Proposition 2.7. Let $\mathcal{I} \subseteq \mathbb{K}$ be a nonzero free $\mathbb{Z}$ -module of rank n . Then

$$(1) \quad \det(\sigma_\alpha(\mathcal{I})) = N(\mathcal{I})^2 N_{\mathbb{K}|\mathbb{Q}}(\alpha) d_{\mathbb{K}}.$$

3. ROTATED D_n -LATTICES

For $n \geq 3$, the n -dimensional lattice D_n in $\mathbb{R}^n$ is described, in its standard form, as $D_n = \{(x_1, \dots, x_n) \in \mathbb{Z}^n; \sum_{i=1}^n x_i \text{ is even}\}$. The set $\beta = \{(-1, -1, 0, \dots, 0), (1, -1, 0, \dots, 0), (0, 1, -1, 0, \dots, 0), \dots, (0, 0, \dots, 1, -1)\}$ is a basis for D_n and $\det(D_n) = 4$ for all n .

As it is known, D_n has a very efficient decoding algorithm and also has the best lattice packing density for $n = 3, 4, 5$ [3].

Using subfields of some cyclotomic fields we have obtained in [9] full diversity rotated D_n -lattices.

Proposition 3.1. [9] Let $\mathbb{K} = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $e_0 = 1$, $e_i = \zeta_{2^r}^i + \zeta_{2^r}^{-i}$ and $n = 2^{r-2}$. If $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is the $\mathbb{Z}$ -module with $\mathbb{Z}$ -basis $\{-2e_0 + 2e_1 - 2e_2 + \dots - 2e_{n-2} + e_{n-1}, -e_{n-1}, e_{n-2}, \dots, (-1)^{i+1}e_{n-1-i}, \dots, e_2, -e_1\}$ and $\alpha = 2 + e_1$, then the lattice $\frac{1}{\sqrt{2^{r-1}}} \sigma_{\alpha}(\mathcal{I}) \subseteq \mathbb{R}^{2^{r-2}}$ is a rotated D_n -lattice and $\mathcal{I} = e_1 \mathcal{O}_{\mathbb{K}}$ is a principal ideal of $\mathcal{O}_{\mathbb{K}}$.

Proposition 3.2. [9] Let $\mathbb{K} = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$, $e_i = \zeta_p^i + \zeta_p^{-i}$ and $n = \frac{p-1}{2}$. If $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is the $\mathbb{Z}$ -module with $\mathbb{Z}$ -basis $\{-e_1 - 2e_2 - \dots - 2e_n, e_1, e_2, \dots, e_{n-1}\}$ and $\alpha = 2 - e_1$, then the lattice $\frac{1}{\sqrt{p}} \sigma_{\alpha}(\mathcal{I}) \subseteq \mathbb{R}^{\frac{p-1}{2}}$ is a rotated D_n -lattice. The $\mathbb{Z}$ -module $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is not an ideal of $\mathcal{O}_{\mathbb{K}}$.

In what follows we consider the compositum of the fields $\mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$ and $\mathbb{Q}(\zeta_p + \zeta_p^{-1})$, p prime, to present new constructions of rotated D_n -lattices. These lattices are obtained via free $\mathbb{Z}$ -modules of rank n which are not ideals.

Remark 3.3. A remark to be used in the next proposition is that if R is a rotation in $\mathbb{R}^n$ and $\mathbf{x} \in R(\mathbb{Z}^n)$ such that $\|\mathbf{x}\|^2$ is even, then $\mathbf{x} \in R(D_n)$. In fact, if we start from $\mathbf{y} \in \mathbb{Z}^n$ such that $\|\mathbf{y}\|^2 = y_1^2 + \dots + y_n^2 = 2a$ for some $a \in \mathbb{N}$, note that $(\sum_{i=1}^n y_i)^2 = \sum_{i=1}^n y_i^2 + 2t = 2a + 2t$, $t \in \mathbb{Z}$, is even. Then, $\sum_{i=1}^n y_i$ is even and $\mathbf{y} \in D_n$. Since a rotation R preserves the norm of a vector, given $\mathbf{x} = R(\mathbf{y})$, $\mathbf{y} \in \mathbb{Z}^n$ such that $\|\mathbf{x}\|^2$ is even, then $\mathbf{x} \in R(D_n)$. Particularly, D_n is the largest even sublattice of $\mathbb{Z}^n$.

Proposition 3.4. Let $\mathbb{K}_1 = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $r \geq 3$, $e_0 = 1$, $e_i = \zeta_{2^r}^i + \zeta_{2^r}^{-i}$, $i \geq 1$, and $n_1 = 2^{r-2}$. Let $\mathbb{K}_2 = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$, $p \geq 5$ odd prime, $b_i = \zeta_p^i + \zeta_p^{-i}$, $i \geq 1$, and $n_2 = \frac{p-1}{2}$. If $\mathbb{K} = \mathbb{K}_1 \mathbb{K}_2$ is the compositum of $\mathbb{K}_1$ and $\mathbb{K}_2$, $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is the $\mathbb{Z}$ -module with $\mathbb{Z}$ -basis

$$\gamma = \{e_0 b_1, e_0 b_2, \dots, e_0 b_{n_2-1}, 2e_0 b_{n_2}, e_1 b_1, \dots, e_1 b_{n_2}, \dots, e_{n_1} b_1, \dots, e_{n_1} b_{n_2}\},$$

$\alpha_1 = (2 - e_1)$, $\alpha_2 = (2 - b_1)$, $\alpha = (2 - e_1)(2 - b_1)$, then the lattice $\frac{1}{\sqrt{2^{r-1}p}} \sigma_{\alpha}(\mathcal{I}) \subseteq \mathbb{R}^n$ is a rotated D_n -lattice.

Proof. From [1] and [5] we have:

$$Tr_{\mathbb{K}_1|\mathbb{Q}}(\alpha_1 e_i e_i) = \begin{cases} 2n_1, & \text{if } i = 0; \\ 4n_1, & \text{if } i \neq 0. \end{cases} \quad Tr_{\mathbb{K}_2|\mathbb{Q}}(\alpha_2 b_j b_j) = \begin{cases} p, & \text{if } j = n_2; \\ 2p, & \text{if } j \neq n_2. \end{cases}$$

Using the fact that $Tr_{\mathbb{K}|\mathbb{Q}}(\alpha_1 e_i b_j \alpha_2 e_k b_l) = Tr_{\mathbb{K}_1|\mathbb{Q}}(\alpha_1 e_i e_k) Tr_{\mathbb{K}_2|\mathbb{Q}}(\alpha_2 b_j b_l)$ we may conclude:

$$Tr_{\mathbb{K}|\mathbb{Q}}(\alpha e_i b_j e_i b_j) = \begin{cases} 2n_1 p, & \text{if } i = 0, j = n_2; \\ 8n_1 p, & \text{if } i \neq 0, j \neq n_2 \\ 4n_1 p, & \text{otherwise.} \end{cases}$$

Considering the Gram matrix G associated to the $\mathbb{Z}$ -basis γ we may assert that the value of the elements of its diagonal are either $Tr_{\mathbb{K}|\mathbb{Q}}(\alpha_2 e_0 b_{n_2} 2e_0 b_{n_2}) = 8n_1 p$ or $Tr_{\mathbb{K}|\mathbb{Q}}(\alpha e_i b_j e_i b_j) \in \{4n_1 p, 8n_1 p\}$ for $i \neq 0$. From [5], we have that $\Lambda_1 = \frac{1}{\sqrt{2^{r-1}p}} \sigma_\alpha(\mathcal{O}_{\mathbb{K}})$ is a rotated $\mathbb{Z}^n$ -lattice. Since $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$, then $\Lambda = \frac{1}{\sqrt{2^{r-1}p}} \sigma_\alpha(\mathcal{I}) \subseteq \Lambda_1$. By Remark 3.3, $\sigma_\alpha(2e_0 b_{n_2} 2e_0 b_{n_2}), \sigma_\alpha(e_i b_j e_i b_j) \in R(D_n)$, where $R(D_n)$ is a rotated version of D_n contained in Λ_1 . Then, $\Lambda \subseteq R(D_n)$. Moreover, $\Lambda = R(D_n)$. In fact, since $d_{\mathbb{K}} = 2^{(r-1)2^{r-2}-1}$ [10], by Proposition 2.7, $\det(\sigma_\alpha(\mathcal{I})) = N(\mathcal{I})^2 N_{\mathbb{K}|\mathbb{Q}}(\alpha) d_{\mathbb{K}} = 2^2 (N_{\mathbb{K}_1|\mathbb{Q}}(\alpha_1))^{n_2} (N_{\mathbb{K}_2|\mathbb{Q}}(\alpha_2))^{n_1} d_{\mathbb{K}_1}^{n_2} d_{\mathbb{K}_2}^{n_1} = 2^2 2^{n_2} p^{n_1} 2^{n_2((r-1)2^{r-2}-1)} p^{n_1 \frac{p-3}{2}} = 4(2^{r-1}p)^{n_1 n_2}$. Therefore, $\Lambda = R(D_n)$ is a rotated D_n -lattice. $\square$

Proposition 3.5. *The $\mathbb{Z}$ -module $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ given in Proposition 3.4 is not an ideal of $\mathcal{O}_{\mathbb{K}}$.*

Proof. Indeed, if $e_0 b_{n_2} \in \mathcal{I}$, then $\mathcal{I} = \mathcal{O}_{\mathbb{K}}$. Since $\left| \frac{\mathcal{O}_{\mathbb{K}}}{\mathcal{I}} \right| = 2$, then $e_0 b_{n_2} \notin \mathcal{I}$. Note that $e_0 b_{n_2-1} e_0 b_1 = e_0 b_{n_2} + e_0 b_{n_2-2}$ is not in $\mathcal{I}$. In fact, if $e_0 b_{n_2-1} e_0 b_1 \in \mathcal{I}$, then $e_0 b_{n_2} = e_0 b_{n_2-1} e_0 b_1 - e_0 b_{n_2-2} \in \mathcal{I}$ and this does not happen. $\square$

Proposition 3.6. *If $\Lambda = \frac{1}{\sqrt{2^{r-1}p}} \sigma_\alpha(\mathcal{I}) \subseteq \mathbb{R}^n$ with α and $\mathcal{I}$ as in Proposition 3.4, then the relative minimum product distance is $d_{p,rel}(\Lambda) = 2^{\frac{-nr+n_2}{2}} p^{\frac{-n+n_1}{2}}$.*

Proof. From [9], $|N_{\mathbb{K}_1|\mathbb{Q}}(b_1)| = 1$. Then, $|N_{\mathbb{K}|\mathbb{Q}}(b_1)| = |(N_{\mathbb{K}_1|\mathbb{Q}}(b_1))^{n_2} N_{\mathbb{K}_2|\mathbb{Q}}(1)^{n_1}| = 1$. By Proposition 2.3, $d_p(\sigma_\alpha(\mathcal{I})) = \sqrt{N_{\mathbb{K}_1|\mathbb{Q}}(\alpha) \min_{0 \neq y \in \mathcal{I}} |N_{\mathbb{K}_1|\mathbb{Q}}(y)|} = \sqrt{2^{n_2} p^{n_1}}$, since $\min_{0 \neq y \in \mathcal{I}} |N(y)| = 1$. The minimum Euclidean norm of D_n is $\sqrt{2}$ what gives the expression for the relative minimum product distance:

$$d_{p,rel}(\Lambda) = \left(\frac{1}{\sqrt{2}} \right)^{n_1 n_2} \left(\frac{1}{\sqrt{2^{r-1}p}} \right)^{n_1 n_2} \sqrt{2^{n_2} p^{n_1}} = 2^{\frac{-n_1 n_2 r + n_2}{2}} p^{\frac{-n_1 n_2 + n_1}{2}}. \quad \square$$

In what follows we consider the compositum of the fields $\mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})$ and $\mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})$, p_1, p_2 distinct prime numbers, to construct rotated D_n -lattices. In this construction we also obtain rotated D_n -lattices via free $\mathbb{Z}$ -modules of rank n which are not ideals.

Proposition 3.7. *Let $\mathbb{K}_1 = \mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})$, $\mathbb{K}_2 = \mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})$, $p_1, p_2 \geq 5$ prime numbers, $p_1 \neq p_2$, $e_i = \zeta_{p_1}^i + \zeta_{p_1}^{-i}$, $b_i = \zeta_{p_2}^i + \zeta_{p_2}^{-i}$. If $\mathbb{K} = \mathbb{K}_1 \mathbb{K}_2$ and $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ is the*

$\mathbb{Z}$ -module with $\mathbb{Z}$ -basis

$$\gamma_1 = \{e_1 b_1, e_1 b_2, \dots, e_1 b_{n_2-1}, e_1 b_{n_2}, e_2 b_1, \dots, e_2 b_{n_2}, \dots, e_{n_1} b_1, \dots, 2e_{n_1} b_{n_2}\},$$

$\alpha_1 = (2 - e_1)$, $\alpha_2 = (2 - b_1)$, $\alpha = (2 - e_1)(2 - b_1)$, then the lattice $\frac{1}{\sqrt{p_1 p_2}} \sigma_\alpha(\mathcal{I}) \subseteq \mathbb{R}^n$ is a rotated D_n -lattice.

Proof. From [5]:

$$Tr_{\mathbb{K}|\mathbb{Q}}(\alpha e_i b_j e_i b_j) = \begin{cases} 2p_1 p_2, & \text{if } i = n_1 \text{ and } j = n_2; \\ 2p_1 p_2, & \text{if either } i \neq n_1 \text{ and } j = n_2 \text{ or } i = n_1 \text{ and } j \neq n_2 \\ 4p_1 p_2, & \text{if } i \neq n_1 \text{ and } j \neq n_2 \end{cases}$$

Using the same ideas of Proposition 3.4, we show that $\frac{1}{\sqrt{p_1 p_2}} \sigma_\alpha(\mathcal{I}) \subseteq R(D_n)$. By Proposition 2.7, $\det(\sigma_\alpha(I)) = 2^2 p_1^{n_2} p_2^{n_1} p_1^{\frac{p_1-1}{3} n_2} p_2^{\frac{p_2-1}{3} n_1} = 4(p_1 p_2)^{n_1 n_2}$. Then, $\det(\Lambda) = 4 = \det(R(D_n))$ and Λ is a rotated D_n -lattice. $\square$

Proposition 3.8. *The $\mathbb{Z}$ -module $I \subseteq \mathcal{O}_{\mathbb{K}}$ given in Proposition 3.7 is not an ideal of $\mathcal{O}_{\mathbb{K}}$.*

Proof. Using the same argument of Proposition 3.5 we show that $e_{n_1} b_{n_2-1} e_{n_1} b_1 \notin I$. $\square$

Proposition 3.9. *If $\Lambda = \frac{1}{\sqrt{2^{r-1}p}} \sigma_\alpha(I) \subseteq \mathbb{R}^n$ as in the Proposition 3.7, then the relative minimum product distance is $d_{p,rel}(\Lambda) = 2^{\frac{-n_1 n_2}{2}} p_1^{\frac{-n+n_2}{2}} p_2^{\frac{-n+n_1}{2}}$.*

Proof. From [9], $|N_{\mathbb{K}_1|\mathbb{Q}}(b_1)| = 1$. So, $|N_{\mathbb{K}|\mathbb{Q}}(b_1)| = 1$. By Proposition 2.3, $d_p(\sigma_\alpha(I)) = \sqrt{N_{\mathbb{K}_1|\mathbb{Q}}(\alpha) \min_{0 \neq y \in I} |N_{\mathbb{K}_1|\mathbb{Q}}(y)|} = \sqrt{2^{n_2} p^{n_1}}$, since $\min_{0 \neq y \in I} |N(y)| = 1$. The minimum Euclidean norm of D_n is $\sqrt{2}$ what gives the expression for the relative minimum product distance:

$$d_{p,rel}(\Lambda) = \left(\frac{1}{\sqrt{2}}\right)^{n_1 n_2} \left(\frac{1}{\sqrt{p_1 p_2}}\right)^{n_1 n_2} \sqrt{p_1^{n_2} p_2^{n_1}} = 2^{\frac{-n_1 n_2}{2}} p_1^{\frac{-n_1 n_2 + n_2}{2}} p_2^{\frac{-n_1 n_2 + n_1}{2}}.$$

$\square$

For $\mathbb{K}_1 = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$, p prime, $\mathbb{K}_2 = \mathbb{Q}(\zeta_p + \zeta_p^{-1})\mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, p prime, and $\mathbb{K}_3 = \mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})\mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})$, p_1, p_2 primes, $p_1 \neq p_2$, if it was possible to construct these rotated D_n -lattices via principal ideals of $\mathcal{O}_{\mathbb{K}_i}$ we would have twice the relative minimum product distances than the ones obtained in our constructions. This fact motivated the discussion on the existence of such constructions via ideals of $\mathcal{O}_{\mathbb{K}_i}$, $i = 1, 2, 3$. It will be shown in the next section that it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}_i}$, $i = 1, 2, 3$.

4. ON THE EXISTENCE OF ROTATED D_n -LATTICES CONSTRUCTED VIA GALOIS EXTENSIONS

From (1), a necessary condition to construct a rotated D_n -lattice, scaled by $\sqrt{c}$ with $c \in \mathbb{Z}$, via fractional ideals of $\mathcal{O}_{\mathbb{K}}$, is the existence of a fractional ideal $\mathcal{I} \subseteq \mathcal{O}_{\mathbb{K}}$ and a totally positive element α such that $4c^n = N_{\mathbb{K}|\mathbb{Q}}(\alpha)N(\mathcal{I})^2d_{\mathbb{K}}$.

Proposition 4.1. [11] *If $\mathcal{P} \subseteq \mathbb{Z}$ is a nonzero prime ideal and $\mathbb{K}|\mathbb{Q}$ is a Galois extension, then $\mathcal{P}\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^g \mathcal{Q}_i^e$, where $\mathcal{Q}_i$'s are distinct nonzero prime ideals of $\mathcal{O}_{\mathbb{K}}$. Moreover, $f(\mathcal{Q}_i|\mathcal{P}) = [\mathcal{O}_{\mathbb{L}}/\mathcal{Q}_i : \mathcal{O}_{\mathbb{K}}/\mathcal{P}\mathcal{O}_{\mathbb{K}}] = f$ and $n = efg$.*

Proposition 4.2. *Let $\mathbb{K}|\mathbb{Q}$ be a Galois extension, $[\mathbb{K} : \mathbb{Q}] = n$, $d_{\mathbb{K}} = 2^z d$, $z \in \mathbb{N}$ and d odd. If $2\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^g \mathcal{Q}_i^e$ and $f = f(\mathcal{Q}_i|2\mathcal{O}_{\mathbb{K}})$ does not divide $2 - z$, then it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}}$.*

Proof. Any ideal $\mathcal{B}$ of $\mathcal{O}_{\mathbb{K}}$ with even norm satisfies $N(\mathcal{B}) = (2^f)^a b$ where $a \geq 1$ and b is odd [11]. Let $\mathcal{I}$ be a fractional ideal of $\mathbb{K}$. By Lemma 2.6, there is $t \in \mathbb{Z}$ such that $t\mathcal{I} = \mathcal{A}$ is an integer ideal of $\mathcal{O}_{\mathbb{K}}$. Then, $N(\mathcal{I}) = N(\mathcal{A})/t^n$. $N(\mathcal{A})$ can be written as $N(\mathcal{A}) = (2^f)^{a_1} b_1$ and so $N(\mathcal{I}) = \frac{(2^f)^{a_1} b_1}{t^n}$, with $a_1 \in \mathbb{N}$ and b_1 odd. Note that $N(\alpha) = N(\alpha\mathcal{O}_{\mathbb{K}}) = (2^f)^{a_2} b_2$ with $a_2 \in \mathbb{N}$, b_2 odd. Let $t = 2^k l$, $k \in \mathbb{N}$ and l odd. Since $n = efg$, for all $c \in \mathbb{Z}$ we have

$$(2) \quad N(\mathcal{I})^2 N(\alpha) d_{\mathbb{K}} = \frac{(2^f)^{2a_1} b_1^2}{t^{2n}} (2^f)^{a_2} b_2 2^z d = 2^{2fa_1 - 2kn + fa_2 + z} (b_1^2 l^{-2n} b_2 d) \neq 4c^n.$$

Indeed, if $c = 2^a b$ with $a \geq 0$ and b odd, then $4c^n = (2^n)^a 2^{2n} b^n$ and the powers of 2 must be equal in (2), therefore $aefg + 2 = an + 2 = f(2a_1 + a_2 - keg) + z$ and this implies that $2 - z = f(2a_1 + a_2 - aeg - keg)$. By hypothesis f does not divide $2 - z$ and then it is impossible to find a fractional ideal $\mathcal{I}$ and an element $\alpha \in \mathcal{O}_{\mathbb{K}}$ satisfying the necessary conditions. $\square$

Remark 4.3. *Let $\mathbb{K} = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$. In [9], we have presented a construction of rotated D_n -lattices via ideals of $\mathcal{O}_{\mathbb{K}}$. Note that $f|(2 - z)$. Indeed, we have $2\mathcal{O}_{\mathbb{K}} = \mathcal{P}^{2^{r-2}}$ where $\mathcal{P} = (2 - (\zeta_{2^r} + \zeta_{2^r}^{-1}))\mathcal{O}_{\mathbb{K}}$. Hence, $f = 1$ and f divides $2 - z$. A natural question to be answered is if the condition $f|(2 - z)$ is sufficient to allow the construction of rotated D_n -lattices via fractional ideals, assuring a kind of Proposition 4.2 reciprocal.*

We will show in Proposition 4.6 that for any Galois extension $\mathbb{K}|\mathbb{Q}$ with discriminant $d_{\mathbb{K}}$ odd it is impossible to construct such rotated D_n -lattices via fractional ideals in $\mathbb{K}$. In particular, this assertion includes the fractional ideals of $\mathbb{Q}(\zeta_p + \zeta_p^{-1})$ and $\mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})\mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})$, p, p_1, p_2 prime numbers, $p \geq 7$, $p_1, p_2 \geq 5$, $p_1 \neq p_2$.

Lemma 4.4. *Let $\mathbb{K}|\mathbb{Q}$ be a Galois extension with odd discriminant. Therefore, $2\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^s \mathcal{Q}_i$ with $\mathcal{Q}_i \neq \mathcal{Q}_j$ prime ideals of $\mathcal{O}_{\mathbb{K}}$.*

Proof. Let $2\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^s \mathcal{Q}_i^e$. From [11], $e > 1$ if, and only if, $2\mathcal{O}_{\mathbb{K}}$ ramifies in $\mathcal{O}_{\mathbb{K}}$. But this happens if, and only if, 2 divides $d_{\mathbb{K}}$, what does not happen. Therefore, $e = 1$. $\square$

In the next lemma we use the Kummer's theorem to factorize ideals as a product of prime ideals.

Lemma 4.5. *If $\mathbb{K} = \mathbb{Q}(\theta)$ is a Galois extension such that $[\mathbb{K} : \mathbb{Q}] \notin \{1, 2, 4\}$ and $2\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^s \mathcal{Q}_i$, $\mathcal{Q}_i \neq \mathcal{Q}_j$, then $f \neq 1$ and $f \neq 2$.*

Proof. Let $m(x) = \min_{\mathbb{Q}}(\theta)$ and $\overline{m(x)}$ the polynomial obtained from $m(x)$ by reduction modulo $\mathbb{Z}_2[x]$. If $f = 1$, $\overline{m(x)}$ is written as the product of irreducible polynomials of degree 1 in $\mathbb{Z}_2[x]$ and there are only two possibilities for these polynomials, namely, $\overline{m_1(x)} = \overline{x}$ and $\overline{m_2(x)} = \overline{x - 1}$. If $\overline{m(x)} = \overline{m_1(x)m_2(x)}$, then $2\mathcal{O}_{\mathbb{K}} = \mathcal{Q}_1\mathcal{Q}_2$ and since $f = 1$, $N(\mathcal{Q}_1) = N(\mathcal{Q}_2) = 2^1 = 2$. So it follows that $N(2\mathcal{O}_{\mathbb{K}}) = 2^2$ and this is possible only if $n = 2$. If $\overline{m(x)} = \overline{m_1(x)}$, then $2\mathcal{O}_{\mathbb{K}} = \mathcal{Q}_1$ and $2^n = N(\mathcal{Q}_1) = 2$ and this is possible only if $n = 1$. If $\overline{m(x)} = \overline{m_2(x)}$, we also get $n = 1$ in a similar way. If $f = 2$, $\overline{m(x)}$ is written as a product of irreducible polynomials of degree 2 in $\mathbb{Z}_2[x]$ and there are only two possibilities for these polynomials: $\overline{m_1(x)} = \overline{x^2}$ and $\overline{m_2(x)} = \overline{x^2 + x + 1}$. If $\overline{m(x)} = \overline{m_1(x)m_2(x)}$, then $2\mathcal{O}_{\mathbb{K}} = \mathcal{Q}_1\mathcal{Q}_2$ and since $f = 2$, then $n = 4$. If $\overline{m(x)} = \overline{m_1(x)}$ or $\overline{m(x)} = \overline{m_2(x)}$ we also get $n = 2$ in a similar way. $\square$

Proposition 4.6. *For any Galois extension $\mathbb{K}|\mathbb{Q}$ of degree $n \notin \{1, 2, 4\}$ and $d_{\mathbb{K}}$ odd, it is impossible to construct a rotated D_n -lattice via a twisted homomorphism applied to a fractional ideal of $\mathcal{O}_{\mathbb{K}}$.*

Proof. By Lemma 4.5, $f \neq \{1, 2\}$. Then f does not divide 2 and the statement follows from Proposition 4.2. $\square$

Corollary 4.7. *If $\mathbb{K}|\mathbb{Q}$ is a Galois extension with conductor m odd, then it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}}$.*

Proof. This result follows immediately from the fact that if the conductor m is odd, $d_{\mathbb{K}}$ is also odd. $\square$

Remark 4.8. *By Corollary 4.7, it is impossible to construct rotated D_3 and D_5 -lattices via fractional ideals of any Galois extension $\mathbb{K} \subseteq \mathbb{Q}(\zeta_m + \zeta_m^{-1})$, m odd.*

In what follows we will show that it is also impossible to construct rotated D_n -lattices via fractional ideals of $\mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})\mathbb{Q}(\zeta_t + \zeta_t^{-1})$, t odd.

The next lemma is presented in [13] considering cyclotomic fields $\mathbb{K}_1 = \mathbb{Q}(\zeta_{2^r})$ and $\mathbb{K}_2 = \mathbb{Q}(\zeta_t)$, t odd. Using the same ideas we may prove the analogous result for $\mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$ and $\mathbb{Q}(\zeta_t + \zeta_t^{-1})$.

Lemma 4.9. *Let $t \in \mathbb{N}$ odd, $\mathbb{K}_1 = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $\mathbb{K}_2 = \mathbb{Q}(\zeta_t + \zeta_t^{-1})$, $[\mathbb{K}_1 : \mathbb{Q}] = n_1$, $[\mathbb{K}_2 : \mathbb{Q}] = n_2$, $\mathbb{K} = \mathbb{K}_1\mathbb{K}_2$. If $2\mathcal{O}_{\mathbb{K}_2} = \mathcal{P}_1\mathcal{P}_2 \cdots \mathcal{P}_r$, $\mathcal{P}_i \neq \mathcal{P}_j$ if $i \neq j$ where $\mathcal{P}_i$ are prime ideals of $\mathcal{O}_{\mathbb{K}_2}$, then $2\mathcal{O}_{\mathbb{K}} = (\mathcal{Q}_1\mathcal{Q}_2 \cdots \mathcal{Q}_r)^{n_1}$, where $\mathcal{Q}_1, \dots, \mathcal{Q}_r$ are prime ideals of $\mathcal{O}_{\mathbb{K}}$ above of $\mathcal{P}_1, \dots, \mathcal{P}_r$, respectively and $f(\mathcal{Q}_i|\mathcal{P}) = f(\mathcal{P}_i|\mathcal{P})$, for all $i = 1, \dots, r$.*

Proof. We have $2\mathcal{O}_{\mathbb{K}_1} = \mathcal{R}^{n_1}$, where $\mathcal{R} = (2 - (\zeta_{2^k} + \zeta_{2^k}^{-1})\mathcal{O}_{\mathbb{K}_1})$ is a prime ideal of $\mathcal{O}_{\mathbb{K}_1}$. Furthermore $2\mathcal{O}_{\mathbb{K}_2} = \mathcal{P}_1\mathcal{P}_2 \cdots \mathcal{P}_r$ a product of distinct prime ideals of $\mathcal{O}_{\mathbb{K}_2}$, with residual degree f . Consider $\mathcal{Q}_1, \dots, \mathcal{Q}_r$ prime ideals of $\mathcal{O}_{\mathbb{K}}$ above of $\mathcal{P}_1, \dots, \mathcal{P}_r$, respectively. We must have that $\mathcal{Q}_i$ is above of $2\mathcal{O}_{\mathbb{K}_1}$ and also above of $\mathcal{R}$ for all $i = 1, \dots, r$. Let $2\mathcal{O}_{\mathbb{K}} = (\mathcal{Q}_1 \cdots \mathcal{Q}_r \mathcal{Q}_{r+1} \cdots \mathcal{Q}_s)^{\bar{e}}$, where $\mathcal{Q}_i$'s are prime ideals of $\mathcal{O}_{\mathbb{K}}$ with residual degree $\bar{f}$. Then, $s\bar{e}\bar{f} = n$. Note that $\bar{e} = e(\mathcal{Q}_i|2) = e(\mathcal{Q}_i|\mathcal{R})e(\mathcal{R}|2) = e(\mathcal{Q}_i|\mathcal{R})n_1$ and $\bar{f} = f(\mathcal{Q}_i|2) = f(\mathcal{Q}_i|\mathcal{P}_i)f(\mathcal{P}_i|2) = f(\mathcal{Q}_i|\mathcal{P}_i)f$. Then, $n = s\bar{e}\bar{f} = se(\mathcal{Q}_i|\mathcal{R})n_1f(\mathcal{Q}_i|\mathcal{P}_i)f \geq rn_1f = n_1n_2$. Therefore, $se(\mathcal{Q}_i|\mathcal{R})n_1f(\mathcal{Q}_i|\mathcal{P}_i)f = rn_1f$, what implies that $se(\mathcal{Q}_i|\mathcal{R})f(\mathcal{Q}_i|\mathcal{P}_i) = r$, with $s \geq r$, $e(\mathcal{Q}_i|\mathcal{R}) \geq 1$, $f(\mathcal{Q}_i|\mathcal{P}_i) \geq 1$. So, $r = s$, $e(\mathcal{Q}_i|\mathcal{R}) = f(\mathcal{Q}_i|\mathcal{P}_i) = 1$, $\bar{e} = n_1$ and $\bar{f} = f$ and then $2\mathcal{O}_{\mathbb{K}} = (\mathcal{Q}_1 \cdots \mathcal{Q}_r)^{n_1}$ and $f = f(\mathcal{Q}_i|\mathcal{P}) = f(\mathcal{P}_i|\mathcal{P})$. $\square$

Proposition 4.10. *Let $\mathbb{K}_1 = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $\mathbb{K}_2 = \mathbb{Q}(\zeta_t + \zeta_t^{-1})$, t odd, $n_1 = [\mathbb{K}_1 : \mathbb{Q}]$ and $n_2 = [\mathbb{K}_2 : \mathbb{Q}] \notin \{1, 2, 4\}$. For $\mathbb{K} = \mathbb{K}_1\mathbb{K}_2$, the compositum of $\mathbb{K}_1$ and $\mathbb{K}_2$, it is impossible to construct rotated D_n -lattices via fractional ideals of $\mathcal{O}_{\mathbb{K}}$.*

Proof. Since $\gcd(d_{\mathbb{K}_1}, d_{\mathbb{K}_2}) = 1$, then $d_{\mathbb{K}} = (d_{\mathbb{K}_1})^{n_2}(d_{\mathbb{K}_2})^{n_1} = (2^{(r-1)2^{r-2}-1})^{n_2}(d_{\mathbb{K}_2})^{n_1}$ with $d_{\mathbb{K}_2}$ odd. Let $2\mathcal{O}_{\mathbb{K}} = \prod_{i=1}^g \mathcal{Q}_i^e$. We will show that $f(2-z) = 2 - ((r-1)2^{r-2} - 1)n_2$. Indeed, $n_1n_2 = efg = n_1fg$ by Proposition 4.9 and hence, $n_2 = fg$. If $f|(2-z)$, then there is $q \in \mathbb{Z}$ such that $fq = 2-z = 2 - ((r-1)2^{r-2} - 1)n_2 = 2 - ((r-1)2^{r-2} - 1)fg$ and $f(q + ((r-1)2^{r-2} - 1)g) = 2$. By hypothesis, $n_2 \notin \{1, 2, 4\}$ and by Proposition 4.5, $f \neq 1$ and $f \neq 2$. Therefore, f does not divide $(2-z)$. $\square$

Remark 4.11. *In Table 1 we compare the relative minimum product distances of rotated D_n -lattices obtained in Propositions 3.1, 3.2, 3.4 and 3.7 considering the number fields $\mathbb{K}_1 = \mathbb{Q}(\zeta_p + \zeta_p^{-1})$, $\mathbb{K}_2 = \mathbb{Q}(\zeta_{2^r} + \zeta_{2^r}^{-1})$, $\mathbb{K}_3 = \mathbb{Q}(\zeta_{2^{r_1}} + \zeta_{2^{r_1}}^{-1})\mathbb{Q}(\zeta_{p_1} + \zeta_{p_1}^{-1})$ and $\mathbb{K}_4 = \mathbb{Q}(\zeta_{p_2} + \zeta_{p_2}^{-1})\mathbb{Q}(\zeta_{p_3} + \zeta_{p_3}^{-1})$.*

In dimensions 8, 128 and 32768 we have considered the Fermat prime numbers 17, 257 and 65537 to construct rotated D_n -lattices via $\mathbb{Z}$ -modules in $\mathcal{O}_{\mathbb{K}_1}$ and via ideals in $\mathcal{O}_{\mathbb{K}_2}$. It is worth noticing that, in these cases, the relative minimum product distances obtained via $\mathbb{Z}$ -modules in $\mathcal{O}_{\mathbb{K}_1}$ are greater than the ones obtained via ideals in $\mathcal{O}_{\mathbb{K}_2}$.

Note also that when both constructions are possible the relative minimum product distances of rotated D_n -lattices obtained via compositum of fields is smaller than the

n	p	r	r_1	p_1	p_2	p_3	$\mathbb{K}_1$	$\mathbb{K}_2$	$\mathbb{K}_3$	$\mathbb{K}_4$
3	7	—	—	—	—	—	0.369646	—	—	—
4	—	4	3	5	—	—	—	0.324210	0.281171	—
5	11	—	—	—	—	—	0.27097	—	—	—
6	13	—	3	7	—	—	0.24285	—	0.219793	—
8	17	5	4	5	—	—	0.20472	0.201311	0.182317	—
10	—	—	3	11	—	—	—	—	0.161122	—
11	23	—	—	—	—	—	0.17003	—	—	—
12	—	—	3	7	—	—	—	—	0.144401	—
14	29	—	—	—	—	—	0.148086	—	—	—
15	31	—	—	—	7	11	0.142402	—	—	0.1380198
16	—	6	5	5	—	—	—	0.133393	0.123452	—
18	37	—	3	19	—	—	0.128512	—	0.1136	—
20	41	—	4	11	—	—	0.121175	—	0.104475	—
128	257	9	—	—	—	—	0.0450746	0.044554	—	—
32768	65537	17	—	—	—	—	0.00276258	0.00276222	—	—

TABLE 1. Relative minimum product distance of rotated D_n -lattices associated to Galois extensions

ones obtained via $\mathbb{K}_1$ and $\mathbb{K}_2$. On the other hand, rotated D_n -lattices constructed via the compositum of fields can be obtained in many dimensions that were not considered before.

5. CONCLUSION

In this paper we have obtained families of full diversity rotated D_n -lattices via $\mathbb{Z}$ -modules in some compositum of Galois fields. It is also shown that for any Galois extension with odd discriminant it is impossible to construct rotated D_n -lattices via fractional ideals of its ring of integers and established a necessary condition to construct rotated D_n -lattices in a Galois extension with even discriminant. As a consequence we can assert that rotated D_n -lattices can not be constructed via fractional ideals in Galois extensions with conductor m odd.

REFERENCES

- [1] A.A. Andrade, C. Alves, T.B Carlos, *Rotated lattices via the cyclotomic field $\mathbb{Q}(\zeta_{2r})$* , International Journal of Applied Mathematics, v.19, n.3, p.321-331, 2006.
- [2] J. Boutros, E. Viterbo, C. Rastello, J.C. Belfiori, *Good lattice constellations for both Rayleigh fading and Gaussian channels*, IEEE Transactions on Information Theory, v.42, n.2, p.502-517, 1996.
- [3] J.H. Conway, N.J.A. Sloane, *Sphere Packings, Lattices and Groups*, Springer-Verlag, 1998.
- [4] E. Bayer-Fluckiger, *Lattices and number fields*, Contemporary Mathematics, v.241, p.69-84, 1999.
- [5] E. Bayer-Fluckiger, F. Oggier, E. Viterbo, *New algebraic constructions of rotated $\mathbb{Z}^n$ -lattice constellations for the Rayleigh fading channel*, IEEE Transactions on Information Theory, v.50, n.4, p.702-714, 2004.

- [6] E. Bayer-Fluckiger, G. Nebe, *On the Euclidean minimum of some real number fields*, Journal de Théorie des Nombres de Bordeaux, 17 no. 2, p. 437-454, 2005.
- [7] E. Bayer-Fluckiger, I. Suarez, *Ideal lattices over totally real number fields and Euclidean minima*, Archiv der Mathematik, v. 86, n.3, p. 217-225, 2006.
- [8] A.L. Flores, J.C. Interlando, T.P. Nóbrega Neto, A.L. Contiero, *A new number field construction of the lattice E_8* , Contributions to Algebra and Geometry, Springer, v. 56, p.1-6, 2012.
- [9] G.C. Jorge, A.J. Ferrari, S.I.R. Costa, *Rotated D_n -lattices*, Journal of Number Theory, v. 132, p. 2397-2406, 2012.
- [10] Lopes, J.O.D. *Discriminants of subfields of $\mathbb{Q}(\zeta_{2^r})$* , Journal of Algebra and its Applications, V.2, p.463-469, 2003.
- [11] P. Samuel, *Algebraic Theory of Numbers*, Paris, Hermann, 1970.
- [12] L.C. Washington, *Introduction to Cyclotomic Fields*, New York, Springer-Verlag, 1982.
- [13] D.A. Marcus, *Number Fields*, New York, Springer-Verlag, 1977.